


# Stereoisomers



# Stereoisomers

## About this Lesson:

In this lesson plan, students will explore compounds called stereoisomers that differ from each other only in the three-dimensional, spatial arrangement of their atoms, but not in the connectivity of their atoms. The connection between stereoisomerism and drug action will also be discussed.

Using Maestro, students will use the 2D Sketcher and 3D Builder tools to learn how to draw different kinds of stereoisomers. Energy minimizations and geometry optimization calculations will be performed using Jaguar. The tile function will be used to compare stereoisomeric relationships between pairs of molecules to determine if they are enantiomers or diastereomers.

## Learning Objectives:

- Define isomer and explain the difference between constitutional isomers and stereoisomers
- Designate configuration of a stereocenter using the Cahn-Ingold-Prelog (CIP) system
- Determine the stereochemical relationship between a pair of molecules

## Lesson Contents:

1. [Setting Up the Maestro Session](#)
2. [Introduction to Stereoisomers](#)
3. [Assigning Configuration using the Cahn-Ingold-Prelog Rules](#)
4. [Enantiomers](#)
5. [Diastereomers](#)
6. [Individual Exercise](#)
7. [Summary, Additional Resources, and References](#)
8. [Glossary of Terms](#)

### Standards Alignment:

- IB Diploma Programme:
  - Models and Representations ([Skill 1](#))
- ACS Guidelines:
  - Understand molecular structure and bonding ([Coursework Guidelines](#))
- AAMC MCAT:
  - Nature of molecules and intermolecular interactions ([5B](#))

### Assessments for Understanding:

The following types of formative assessments are embedded in this lesson:


- Assessment of student understanding through discussion of warm-up questions and filling in any knowledge gaps about constitutional isomers
- Visual assessment of student-generated enantiomers and diastereomers

### Associated Documentation Pages:


### Warm-Up Questions:

Watch the [Khan Academy videos](#) on stereochemistry.

- 1) What is the difference between constitutional isomers and stereoisomers?
- 2) Label the following pair of molecules as either constitutional isomers or stereoisomers. Both the skeletal structure and bond line structure of each molecule have been provided.



- 3) How many chiral center(s) does the following molecule have?



**Need help?** Contact us at [teaching@schrodinger.com](mailto:teaching@schrodinger.com)

# 1. Setting Up the Maestro Session

At the start of the Maestro session, it is essential to 1) check your mouse actions, 2) change the file path to the Working Directory for this lesson, and 3) save your project file. The working directory indicated in this section contains the files necessary to complete this lesson. If you do not set the appropriate working directory, you will be unable to run any calculations.

1. Launch the Virtual Cluster





Figure 1-1. Virtual workstation login page.



2. Double-click the **course-data** folder on the desktop

Figure 1-2. Course-data folder on the desktop.



3. Right-click the Stereoisomers folder and select **Copy to > Desktop**

Figure 1-3. Copy the lesson folder to the Desktop.



4. Double-click the Maestro icon on the desktop






Figure 1-4. Change the mouse actions.

5. Check your mouse actions.
  - o Go to **Workspace > Customize Mouse Actions**
  - o *Note:* This lesson was made with a three-button mouse with a scroll wheel, but a trackpad can still be used
  - o **Trackpad keys:**
    - **Up/Down trackpad** = Zoom In/Out
    - **Option** = Rotate
    - **Control** = Translate



6. Go to File > Change Working Directory


Figure 1-5. Change Working Directory option.



7. Navigate to Desktop > Stereoisomers folder and click **Choose**


Pre-generated input and results files are included for running jobs or examining output

Figure 1-6. Change Working Directory panel.



8. Go to File > Save Project As

Figure 1-7. Save Project option.



9. Change the *File name* to SN2\_lesson, click Save
- The project is now named SN2\_lesson.prj

Figure 1-8. Save Project panel.

## 2. Introduction to Stereoisomers

Isomers are molecules that have the same molecular formula but have differing structural arrangements in space. Though they are made up of the same components, distinct isomers do not necessarily bear resemblance to each other both visually and in terms of their properties. Isomers can be broken into two types: constitutional isomers and stereoisomers.


**Constitutional isomers** are when two molecules share the same chemical formula but differ in structural arrangement. For example, butane (or *n*-butane) and 2-methylpropane (or isobutane) both have a molecular formula of  $C_4H_{10}$ , but they differ in connectivity. *N*-butane is more linear whereas isobutane is more branched, yet they have the same number of carbon and hydrogen atoms in each isomer.

Constitutional isomers of  $C_4H_{10}$




**Figure 2-1.** Constitutional isomers of  $C_4H_{10}$

**Stereoisomers** have an identical molecular formula and atomic sequence but differ in how they are spatially arranged. *Cis-trans* stereoisomerism can be used to denote disubstituted cycloalkanes. For example, *cis*-1,2-dimethylcyclohexane and *trans*-1,2-dimethylcyclohexane both have the same molecular formula of  $C_8H_{16}$  as well as the same connectivity of atoms, but the spatial arrangement of the methyl groups differ. The *cis* stereoisomer shows the two methyl groups on the same side of the ring, while the *trans* stereoisomer shows the two groups on opposite sides of the ring.



**Figure 2-2.** *Cis-Trans* isomers of  $C_8H_{16}$

Furthermore, the terms *cis* and *trans* are also used to describe stereoisomerism in double bonds. For example, *cis*-2-butene and *trans*-2-butene both have the same molecular formula of  $C_4H_8$  as well as the same connectivity of atoms, but the spatial arrangement of the methyl groups differ. The *cis* stereoisomer shows the groups on the same side of the double bond, while the *trans* stereoisomer exhibits groups on opposite sides of the double bond. These two molecules have two different physical properties because the double bond does not have free rotation like single bonds do at room temperature.



**Figure 2-3.** *Cis-Trans* isomers of  $C_4H_8$

The last type of stereoisomer delves into a specific property called **chirality**. Chiral objects are **not superimposable** on their mirror images. The most common source of molecular chirality is the presence of a chiral center which is a carbon atom with four unique, or different, substituents attached to it. The tetrahedral  $sp^3$ -hybridized carbon atom in bromochloroethanol shown below is chiral because it contains four different substituents:  $-OH$ ,  $-Cl$ ,  $-Br$ , and  $-CH_3$ . We will go into much more detail on how to assign chirality and how to label stereoisomeric relationships of two chiral molecules later in this lesson.




Figure 2-4. Stereoisomers of  $C_2H_4OCIBr$


### 3. Assigning Configuration using the Cahn-Ingold-Prelog (CIP) Rules

In order to determine the specific configuration of a stereocenter, a system developed by three chemists named Cahn, Ingold, and Prelog, was developed.


- **Step 1:** Identify the chiral center and rank the substituents around the chiral center. The higher that atomic number, the higher the priority.
  - If priority can't be assigned by the direct atoms bonded to the chiral center, look to the next set of atoms. Priority is assigned at the first point of difference
  - Atoms in a double or triple bond are considered to be bonded to an equivalent number of similar atoms by single bonds
- **Step 2:** Orient or rotate the molecule so the lowest ranked atom or group points towards the back (on a dashed bond).
- **Step 3:** Read the 3 highest priority groups.
  - If the 3 groups are read clockwise, the configuration is (*R*).
  - If the 3 groups are read counterclockwise, the configuration is (*S*).




**Practice #1.** Assign configuration of (*R*) or (*S*) to the following chiral molecule.




- **Step 1:** After looking at the periodic table, Br has the highest atomic number of 35, then Cl has the second highest atomic number of 17, then O has the third highest atomic number of 8, and C has the lowest atomic number of 6.



- **Step 2:** Since C has the lowest atomic number making it the lowest priority group, we have to rotate the methyl group to the back (on a dashed bond). Then we can relabel our priority groups.



- **Step 3:** The sequence 1 > 2 > 3 follows a clockwise configuration, so we can assign the chiral center as (*R*).




## Computational Exercise #1: Geometry Optimization using Maestro

Even though we draw bonds coming towards us as wedged bonds and bonds going away from us as dashed bonds, sometimes it is still difficult to visualize the three-dimensionality

of a molecule when it is drawn on a flat piece of paper. Now we are going to learn how to build a three-dimensional structure of a chiral molecule, optimize its geometry using Maestro, and use the CIP rules to assign its stereochemical configuration. Visualizing the molecular structure can be done in three different ways:

- 1) Through the 2D Sketcher
- 2) Through the 3D Builder
- 3) Or, by importing pre-built molecules

We will be using the 2D Sketcher tool in Maestro.



### Part 1. Build (R)-1-bromo-1-chloroethane using 2D Sketcher

Before optimizing any molecular geometry, you will need a starting molecule in your workspace. Let's draw the structure of 1-bromo-1-chloroethane using the 2D sketcher.

- 4) Go to **Edit > 2D Sketcher**

Figure 3-1. Opening 2D Sketcher




Figure 3-2. Drawing 1-bromo-1-chloroethane.

- 5) **Draw** 1-bromo-1-chloroethane exactly how it is shown in Figure 3-2.
  - a) Go to the selection bar on the left
  - b) Create a methyl group by choosing the “draw line” and **clicking** in the 2D Workspace
  - c) Drag rightwards from the methyl group to add another carbon
  - d) Create a wedge bond by choosing the wedge bond and dragging down from the central carbon
  - e) Put your cursor over the end of the wedge and type Cl to add a chlorine
  - f) Create a dashed bond by choosing the dashed bond and dragging to the left from the central carbon
  - g) Put your cursor over the end of the wedge and type Br to add a bromine
  - h) Once again click on the “draw line” and click on the “H”. Drag upwards from the central carbon
  - i) Try to replicate the sketch shown in the figure as closely as possible
- 6) Click on **Save as New** and for *Input Entry Title* write

**1-bromo-1-chloroethane**. Click **OK**.




Figure 3-3. Clicking the Tasks button in the upper righthand corner.

## Part 2. Optimize the Geometry using Quantum Mechanic DFT

Now let's optimize the molecule at the quantum mechanical level.

- 7) With the 1-bromo-1-chloroethane entry selected and included, go to **Tasks > Optimization** using Jaguar
  - a) The Jaguar Optimization panel opens
- 8) Use structures from **Project Table (1 selected entry)**




Figure 3-4. Opening the Jaguar Optimization panel.

Various settings can be altered depending on the specific use case. We will only adjust the *Input* tab for this example.




Figure 3-5. Loading the structure and setting QM parameters in the Jaguar Optimization panel

- 9) For *Theory*, select B3LYP-D3
- 10) For *Basis set*, select LACVP\*\*

A few additional comments about preparing for an optimization calculation:

- If you hover the mouse over the basis set in the table, you can see the number of basis functions associated with the basis set. This is useful to know since the quality of the basis set *usually* improves as the number of functions increases, noting again the trade-off between quality and computational expense

- Always make sure the charge and the spin multiplicity are correct (in this case, 1-bromo-1-chloroethane is a neutral, singlet, so charge = 0 and spin multiplicity = 1). Multiplicity is defined as  $2S + 1$ , where  $S$  is the total orbital spin of the molecule. For instance, a system with one unpaired electron is a doublet (multiplicity = 2), since the total orbital spin  $S = \frac{1}{2}$ .
- Use the *Atom-Level Settings* button to define per-atom basis sets
- Constraints can be defined on the *Optimization* tab
- Properties, such as atomic charges, vibrational frequencies, surfaces and more can be requested on the *Properties* tab
- Solvent can be defined via several implicit solvation models on the *Solvation* tab. Note that this example is a gas-phase geometry optimization
- Read more about geometry optimization with Jaguar [here](#)




Figure 3-6. Naming and running the job.

- 11) Change the *Job name* to **1-bromo-1-chloroethane\_opt**
  - a) Usually we incorporate stars (\*) and pluses (+) into file names with S and P, respectively
- 12) Adjust the job settings by clicking the gear button to the right of the Job name
  - o Choose cpu-preemptible (400) as your host with 1 processor
- 13) Click **Run**




Figure 3-7. Banner that shows the job is finished – Result has been incorporated.




Figure 3-8. The output molecule of 1-bromo-1-chloroethane.




Figure 3-9. Viewing and styling 1-bromo-1-chloroethane.

When the job finishes, a banner will appear indicating that the result has been incorporated. At this time, the molecule with the optimized geometry is now shown in the workspace

- 14) A new entry group is added to the entry list titled 1-bromo-1-chloroethane\_opt\_B3LYP-D3\_6-31GSS.011 (1). Select and include the molecule titled 1-bromo-1-chloroethane in this subgroup. This is your optimized structure.

- 15) Change the representation to ball-and-stick by clicking on the **Style** menu and choosing **Apply ball-and-stick representation**
- 16) Change the color of the atoms by clicking on the **Style > Color Atoms > Element + Custom Ligand**




Figure 3-10. Changing atom colors for 1-bromo-1-chloroethane.




Figure 3-11. Labeling atoms.

- 17) To display atom labels, click **Style > Apply Labels** then click the drop down arrow on the right side
- 18) Click **Edit Custom Labeling**
- 19) Checkbox **Element** and press **OK**
- 20) Go back to **Style > Apply Labels > Custom Label** and now each atom in your workspace should be labeled

*Note:* You can toggle the atom labels on and off by clicking Annotations label button in the Workspace toggle on the bottom right corner




Figure 3-12. Toggling labels on.



Figure 3-13. Toggling labels off.



**Pause & Think #1.** Take a screenshot of your optimized structure of (R)-1-bromo-1-chloroethane including atom and stereochemical labels. Choose the ball-and-stick representation styling tool.




Figure 3-14. Determining where the chiral center is.

### Part 3. Assign stereochemical configuration using CIP rules

Now that we optimized the geometry of 1-bromo-1-chloroethane, let's go through the CIP rules to determine its stereochemical configuration.

- 21) Determine which carbon atom is the chiral center. Remember that a chiral carbon has 4 unique, or different, substituents around it




Figure 3-15. Labeling priorities of each substituent around the chiral center.

- 22) Label the priorities of each atom surrounding the chiral center based on atomic number (this can be found from a periodic table). Feel free to take note of these priorities on a separate piece of paper




Figure 3-16. Rotating the molecule to the lowest priority group is facing the back.

23) Rotate the molecule so the lowest priority group is facing the back





Figure 3-17. Assigning configuration based on

24) Relabel priorities and determine whether 1 > 2 > 3 is clockwise (*R*) or counterclockwise (*S*)

25) Since groups 1 > 2 > 3 are going clockwise, this particular isomer is (*R*)-1-bromo-1-chloroethane

either clockwise or counterclockwise sequence of the first 3 priority groups.



- 26) To double-check your answer and display stereochemistry labels, click **Style > Apply Labels** then click the drop down arrow on the right side
- 27) Click **Stereochemistry + Charge** and you will find that indeed, this stereocenter is (*R*)
- 28) Double-click the entry name in the entry list and change the name to R-1-bromo-1-chloroethane.

Figure 3-18. Applying the Stereochemistry + Charge Label to the molecule.

## 4. Enantiomers

A pair of compounds with the same atom connectivity, molecular formula, and only 1 chiral center can either be identical or enantiomers with one another. If they are identical, they would have the same exact stereochemical configuration. If they have completely opposite configurations (i.e. one molecule is (*R*) while the other is (*S*)), then they would be enantiomers. Enantiomers are stereoisomers that are nonsuperimposable, mirror images of one another and exhibit identical physical properties. When exposed to plane-polarized light, however, enantiomers exhibit different behavior. An enantiomer that rotates plane-polarized light in the positive direction is called dextrorotatory [(+), or d-], while the enantiomer that rotates the light in the negative direction is called levorotatory [(-), or l-]. When both d- and l- isomers are present in equal amounts, the mixture is called a **racemic mixture**. To learn more about the optical activity of enantiomers, you can read more about it [here](#).

To better understand and visualize a pair of enantiomers, let's draw the enantiomer of (*R*)-1-bromo-1-chloroethane in Maestro.

### Computational Exercise #2: Compare Multiple Molecules by Tiling




Figure 4-1. Duplicating *R*-1-bromo-1-chloroethane in the Entry List.

1. **Right-click** the entry *R*-1-bromo-1-chloroethane
2. Click **Duplicate > In Place** and you will see a new entry appear in the entry list

ENTRY LIST		
Row	In	Title
1	<input type="radio"/>	1-bromo-1-chloroethane
		▼ 1-bromo-1-chloroethane...
2	<input type="radio"/>	R-1-bromo-1-chloroetha...
	1	▼ 1-bromo-1-chloroethane...
3	<input checked="" type="radio"/>	R-1-bromo-1-chloroetha...

Figure 4-2. Including and selecting the new entry.

3. Include and select the new entry




Figure 4-3. Changing the wedged bond to a dashed bond and updating entry.

4. Go to **Edit > 2D Sketcher**
5. Change the wedged C-Br bond to a dashed C-Br bond
6. Click **Update Entry**




Figure 4-5. Applying the Stereochemistry + Charge label to the molecule.

7. Change the representation to ball-and-stick by clicking on the **Style** menu and choosing **Apply ball-and-stick representation**
8. Change the color of the atoms by clicking on the **Style > Color Atoms > Element + Custom Ligand**
9. To display stereochemistry labels, click **Style > Apply Labels > Stereochemistry + Charge Number**

The screenshot shows the 'ENTRY LIST' table with the following data:

Row	In	Title
1	<input type="radio"/>	1-bromo-1-chloroethane
		▼ 1-bromo-1-chloroethane...
2	<input type="radio"/>	R-1-bromo-1-chloroetha...
		▼ 1-bromo-1-chloroethane...
3	<input checked="" type="radio"/>	S-1-bromo-1-chloroethane

Figure 4-6. Changing the Entry Name.

10. Change the Entry name to S-1-bromo-1-chloroethane by double-clicking on the name.

Row	In	Title
1	<input type="radio"/>	1-bromo-1-chloroethane
1	<input checked="" type="radio"/>	1-bromo-1-chloroethane...
2	<input checked="" type="radio"/>	R-1-bromo-1-chloroetha...
1	<input checked="" type="radio"/>	1-bromo-1-chloroethane...
3	<input checked="" type="radio"/>	S-1-bromo-1-chloroetha...

Figure 4-7. Including both molecules in the Entry List.

Now let's view our molecules side by side.

11. Simultaneously **include** (control + click or command + click) both the R-1-bromo-1-chloroethane and the S-1-bromo-1-chloroethane




Figure 4-8. Opening the Workspace Configuration Panel

12. Click the + symbol in the bottom left-hand corner to show workspace configuration panel
13. Click **Tile** under the Workspace Layout




Figure 4-9. Tiling the two enantiomers side-by-side.




Figure 4-10. Changing the background color to white.


14. Change the background color to white by clicking on the + symbol in the bottom right-hand corner



**Pause & Think #2.** Take a screenshot of the tiled enantiomers for 1-bromo-1-chloroethane including atom and stereochemical labels. Choose the ball-and-stick representation styling tool.

## 5. Diastereomers


Diastereomers are stereoisomers that are not mirror images of one another. The difference between enantiomers and diastereomers is more apparent when we compare compounds with more than one chiral center. For example, let's take a look at the structure of 2-methylcyclohexanol.



**Figure 5-1.** Structure of 2-methylcyclohexanol

This molecule contains two chiral centers where each one can either have the (*R*) configuration or the (*S*) configuration. A molecule can have  $2^n$  number of possible

stereoisomers where  $n$  is the number of chiral centers. Since 2-methylcyclohexanol has 2 chiral centers, it has a total of  $2^2 = 4$  possible stereoisomers, as shown below.



**Figure 5-2.** The possible stereoisomers of 2-methylcyclohexanol

Stereoisomers #1 and #2 are enantiomers because both chiral centers completely inverted to the opposite configuration. The same goes for stereoisomers #3 and #4 where the two chiral centers in stereoisomer #3 were (1*R*, 2*R*) and they both inverted to the opposite configuration of (1*S*, 2*S*) in stereoisomer #4. Now if we compare stereoisomer #1 to stereoisomer #3, notice that one chiral center stayed the same (1*R*) whereas the second chiral center inverted from (2*S*) to (2*R*). Since only one chiral center inverted, these two isomers are no longer mirror images from one another, making them diastereomers. And the same goes for stereoisomers #2 and #4 where they are both diastereomers as well.

As previously mentioned, *cis-trans* isomers of alkenes are also considered to be diastereomers because they too are not mirror images of one another. Let's build a few diastereomeric alkenes to visually compare their differences.

### **Computational Exercise #3: Build Diastereomeric Alkenes**




Figure 5-3. Open 2D Sketcher.

In order to visualize the molecules and compare isomers in three dimensions, we first have to draw models of the molecules. This can be done either through the 2D Sketcher, the 3D builder, or by importing pre-built molecules. We will be using the 2D Sketcher tool:

1. Go to **Edit** → **2D Sketcher**




Figure 5-4. Drawing and saving cis-2-pentene.

2. **Draw** cis-2-pentene
  - a. Go to the selection bar on the left
  - b. Create a methyl group by choosing the “C” or the draw line and **clicking** in the 2D Workspace
  - c. Drag from the methyl group to add more
  - d. Click on the bond between methyl groups to make it a double bond
  - e. Try to replicate the sketch shown in the figure as closely as possible
3. Click on **Save as New** and for *Input Entry Title* write **cis-2-pentene**. Click **OK**.




Figure 5-5. Drawing and saving trans-2-pentene.

4. **Draw trans-2-pentene**
  - a. Reference the instructions for cis-2-pentene in step 1
  - b. Try to replicate the sketch shown in the figure as closely as possible
5. Click on **Save as New** and for *Input Entry Title* write **trans-2-pentene**. Click **OK**

The screenshot shows the 'ENTRY LIST' panel with a table of molecules. The 'cis-2-pentene' and 'trans-2-pentene' entries are selected, indicated by blue circles in the 'In' column.

Row	In	Title
1	<input checked="" type="checkbox"/>	cis-2-pentene
2	<input checked="" type="checkbox"/>	trans-2-pentene
3	<input type="checkbox"/>	cis-1,2-dichlorocyclohexane
4	<input type="checkbox"/>	trans-1,2-dichlorocyclohexane

Figure 5-6. Including the cis/trans molecules

Oftentimes, viewing multiple molecules simultaneously can be important for comparing and contrasting them. We will show our cis and trans isomer molecules side by side so that we can easily see how they differ.




Figure 5-7. Tiling the cis/trans entries.

6. Simultaneously **include** (control + click or command + click) both the cis-2-pentene and the trans-2-pentene
7. Show workspace configuration panel → Workspace Layout → Tile




Figure 5-8. Applying ball-and-stick style to *cis/trans* entries.




Figure 5-9. Viewing 2-pentene isomers.

8. Change the representation to ball-and-stick by clicking on the **Style** menu and choosing **Apply ball-and-stick representation**

9. Feel free to pan around, zoom in, and look at your molecules
10. Change the background to white and take a screenshot of your molecules to put in the Stereoisomers worksheet.

For 2-pentene, notice that the *cis* configuration shows the methyl and the ethyl substituents on the same side of the double bond. The *trans* configuration has the methyl and ethyl on opposite sides of the double bond.






**Pause & Think #3.** Take a screenshot of the tiled *cis* and *trans* diastereomers of 2-pentene including atom and stereochemical labels. Choose the ball-and-stick representation styling tool.





## 6. Individual Exercise


For each pair of molecules:

- 1) Build the molecules using 2D sketcher in Maestro and Optimize their geometries by running Optimization Jaguar calculations. Attach screenshots of your optimized molecules tiled next to each other in the worksheet
- 2) Determine the stereochemical configuration using CIP rules
- 3) Determine whether the pair of molecules are identical, enantiomers, or diastereomers

Here is an example:

 <p><b>Molecule A</b></p> <p><b>Molecule B</b></p>	<p>1)</p> <div style="display: flex; justify-content: space-around;"> <div style="text-align: center;"> <p>Title: Molecule B</p>  </div> <div style="text-align: center;"> <p>Title: Molecule A</p>  </div> </div> <p>2) Molecule A: (1<i>S</i>, 3<i>R</i>)-1-chloro-3-methylcyclopentane Molecule B: (1<i>R</i>, 2<i>R</i>)-1-chloro-3-methylcyclopentane</p> <p>3) Diastereomers because one chiral center inverted while the other chiral center stayed the same</p>
---	--

<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p><b>Molecule A</b></p> </div> <div style="text-align: center;">  <p><b>Molecule B</b></p> </div> </div>	<p>1)</p> <p>2)</p> <p>3)</p>
<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p><b>Molecule A</b></p> </div> <div style="text-align: center;">  <p><b>Molecule B</b></p> </div> </div>	<p>1)</p> <p>2)</p> <p>3)</p>



## 7. Summary, Additional Resources, and References

Using Maestro, students learned how to use Jaguar's AutoTS panel to perform geometry optimization calculations of the reactants and products of an SN2 reaction, determine which bonds are breaking and forming, establish correspondence between atoms in the reactants and the products, and generate a transition state structure. Lastly, they learned how to render a potential energy surface diagram to show the activation energy barrier.

### For further learning:

For some related practice, proceed to explore other relevant tutorials:

- [Introduction to Multistage Quantum Mechanical Workflows](#)

### For further reading:

- [Introduction to Computational Chemistry, 3rd Edition](#)

- [Essentials of Computational Chemistry: Theories and Models, 2nd Edition](#)
- See the Jaguar help [documentation](#)

## 8. Glossary of Terms

Entry List - a simplified view of the Project Table that allows you to perform basic operations such as selection and inclusion

Included - the entry is represented in the Workspace, the circle in the In column is blue

Project Table - displays the contents of a project and is also an interface for performing operations on selected entries, viewing properties, and organizing structures and data

Recent actions - This is a list of your recent actions, which you can use to reopen a panel, displayed below the Browse row. (Right-click to delete.)

Scratch Project - a temporary project in which work is not saved. Closing a scratch project removes all current work and begins a new scratch project

Selected - (1) the atoms are chosen in the Workspace. These atoms are referred to as "the selection" or "the atom selection". Workspace operations are performed on the selected atoms. (2) The entry is chosen in the Entry List (and Project Table) and the row for the entry is highlighted. Project operations are performed on all selected entries

Working Directory - the location that files are saved

Workspace - the 3D display area in the center of the main window, where molecular structures are displayed